



IMMUNE GLOBULIN INFUSION REFERRAL FORM

Phone (888) 383-4852 • Fax: (888) 383-8430

Today's Date

- CURRENT PATIENT
- NEW PATIENT

Patient Name _____ SS# _____ DOB _____
 Height _____ Weight _____ Address _____ Apt # _____
 Male Female City _____ State _____ Zip _____
 Daytime Tel _____ Cell _____ Email _____
 Ship to Patient at Home Work OR Patient will pick up at Physician Office Pharmacy Date Needed _____
 Medical History: Cardiac Disease Diabetes Renal Dysfunction IgA Deficient
 Allergies _____ Comorbidities _____
 Current Medications (if necessary, please fax a complete list) _____

Diagnosis: D80.0 Hereditary Hypogammaglobulinemia D81.5 Immune Deficiency with Increased IgM
 D83.9 Common Variable Immunodeficiency, unspecified D82.0 Wiskott Aldrich Syndrome
 D81.9 Combined Immunodeficiency, unspecified Other _____

Primary Insurance Company: _____ Policy#: _____ Group ID#: _____
 Policyholder Name: _____ Rx BIN#: _____ Rx PCN#: _____
 Policyholder DOB: _____ Relation to Patient: _____ Type: HMO PPO PBM Medicare Medicaid
 Secondary Insurance Company: _____ Policy#: _____ Group ID#: _____
 Policyholder Name: _____ Rx BIN#: _____ Rx PCN#: _____
 Policyholder DOB: _____ Relation to Patient: _____ Type: HMO PPO PBM Medicare Medicaid

Prescriber's Name _____ Office Contact _____
 Street Address _____ Suite# _____ City _____ State _____ Zip _____
 Tel _____ Fax _____ Email _____
 License# _____ NPI# _____ UPIN# _____ DEA# _____

PRESCRIPTION PLEASE ATTACH COPIES OF PATIENT'S INSURANCE CARDS AND HISTORY OF IG LABS

Is this the first dose? Yes No Date of last infusion _____
 If no, list product _____ Next dose due _____

ADMINISTER IVIG:
 _____ gm/kg or _____ grams every _____ wk(s) for _____ cycle(s)
 Rate per manufacturer recommendation Other _____
 Other _____

ADMINISTER SQIg:
 IG _____ grams each month given as _____ doses IG _____ grams _____ times per month.
 Administer SQIg using _____ sites at a time. Repeat _____ week(s). OK to round dose to the nearest vial size. Refill x 1 year.
 Other _____

PRE-MEDICATIONS
 Diphenhydramine (Benadryl) 25-50 mg orally before infusion Other _____
 Acetaminophen (Tylenol) 325-650 mg orally before infusion

ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS CARE PROTOCOL
 Adults or Children greater than 66 pounds or 30 kg:
 • For mild reaction: give Diphenhydramine 50 mg orally, IM or IV and decrease the rate of infusion.
 • For moderate reaction: stop infusion, give Diphenhydramine 50mg, orally, IM or IV and contact physician
 • For Severe reaction w/breathing problem: stop infusion, call 911, give Epinephrine 0.3mg/0.3ml subcutaneously, Diphenhydramine 50 mg IV or IM. Begin NS 0.9% NaCl 500ml IV at a rate of 100-150ml/hr and contact physician.
 Note: **Dosage adjustment necessary for children less than 30kg or 66 pounds:** Diphenhydramine 1.25mg/kg orally, IM or IV with a maximum of 50mg. If Epinephrine is needed 0.15mg/0.15ml 1:1000 subcutaneously

Nursing: Start PIV as required for administration and nurse to administer infusion in home. Access: Peripheral PICC
 Flushing: INS Standards Care Protocol (Heparin, 0.9% NaCl, D5W) Port
 Lab Order(s): _____ Other _____

By signing this form and utilizing our services, you are authorizing the pharmacy and it's employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature (signature required. NO STAMPS) _____ Date _____

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.