

IMMUNE GLOBULIN NEUROLOGY REFERRAL FORM

Today's Date	

Toll-Free (888) 404-6026 • Fax: (567) 806-3216

CURRENT PATIENT
NEW PATIENT

Patient Name	SS#	DOB
Height Weight Address Male Female City Daytime Tel Cell F		Apt #
☐ Male ☐ Female City	State	Zip
Daytime Tel	Email	
Ship to Patient at 🔲 Home 🔲 Work OR Patient will pick up at 🔲	Physician Office 🔲 Pha	rmacy Date Needed
Medical History: 🔲 Cardiac Disease 🔲 Diabetes 🔲 Renal Dy	rsfunction 🔲 IgA Defici	ent
Allergies	Comorbidities	
Current Medications (if necessary, please fax a complete list)		
Diagnosis:	IDP) ☐ M36.0 I ☐ G25.82 ☐ G35 Mu	Lambert-Eaton Syndrome, unspecified Dermatomyositis Stiff-Person Syndrome ultiple Sclerosis (MS)
Primary Insurance Company:		
Policyholder Name:	Rx BIN#:	Rx PCN#:
Policyholder DOB:Relation to Patient:	Туре: 🗖 НМО 🖣	□PPO □PBM □Medicare □Medicaid
Secondary Insurance Company:	Policy#:	Group ID#:
Policyholder Name:		
Policyholder DOB:Relation to Patient:	Type: □HMO □	PPO PBM Medicare Medicaid
Prescriber's Name	Office Contact	
Street Address Suite#_		-
Tel Fax	Email	
License# NPI#	UPIN#	DEA#
		ICE CARDS AND HISTORY OF IG LABS
	S OF PATIENT'S INSURAN	ICE CARDS AND HISTORY OF IG LABS
PRESCRIPTION PLEASE ATTACH COPIE	S OF PATIENT'S INSURAN Date of last infusion	
PRESCRIPTION PLEASE ATTACH COPIE Is this the first dose? \(\text{Yes} \) No If no, list product	S OF PATIENT'S INSURAN Date of last infusion	
PRESCRIPTION Is this the first dose? Yes No If no, list product ADMINISTER IVIG:	Date of last infusion Next dose due	
PRESCRIPTION PLEASE ATTACH COPIE Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then	Date of last infusion Next dose due grams every wk(s)) for cycle(s)
PRESCRIPTION Is this the first dose? □ Yes □ No If no, list product ADMINISTER IVIG: □ 2 grams/kg over days, as a loading dose, then; □ gm/kg or grams;	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
PRESCRIPTION PLEASE ATTACH COPIE Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
PRESCRIPTION Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
PRESCRIPTION Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
PRESCRIPTION Is this the first dose? ☐ Yes ☐ No If no, list product	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
PRESCRIPTION Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion	Date of last infusion Next dose due wk(s) every) for cycle(s) wk(s) for cycle(s)
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS OF INSTANDARDS OF INSTANDAR	Date of last infusion Next dose due grams every wk(s) every) for cycle(s) wk(s) for cycle(s)
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS Of Adults or Children greater than 66 pounds or 30 kg:	Date of last infusion Next dose due wk(s) every care PROTOCOL) for cycle(s) cycle(s)
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS OF INSTANDARDS OF INSTANDAR	Date of last infusion Next dose due wk(s) every wk(s) every and decrease the rate of i) for cycle(s) cycle(s) nfusion.
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS (Adults or Children greater than 66 pounds or 30 kg: • For mild reaction: give Diphenhydramine 50 mg orally, IM or IV • For moderate reaction: stop infusion, give Diphenhydramine 50m • For Severe reaction w/breathing problem: stop infusion, call 911	Date of last infusion Next dose due wk(s) every wk(s) every and decrease the rate of ing, orally, IM or IV and co, give Epinephrine 0.3mg/	of for cycle(s) cycle(s) on fusion. Infusion. In
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS Of Adults or Children greater than 66 pounds or 30 kg: • For mild reaction: give Diphenhydramine 50 mg orally, IM or IV • For moderate reaction: stop infusion, give Diphenhydramine 50mg	Date of last infusion Next dose due wk(s) every wk(s) every and decrease the rate of ing, orally, IM or IV and co, give Epinephrine 0.3mg/	of for cycle(s) cycle(s) on fusion. Infusion. In
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS (Adults or Children greater than 66 pounds or 30 kg: • For mild reaction: give Diphenhydramine 50 mg orally, IM or IV • For moderate reaction: stop infusion, give Diphenhydramine 50m • For Severe reaction w/breathing problem: stop infusion, call 911	Date of last infusion Next dose due grams every wk(s) every CARE PROTOCOL and decrease the rate of ing, orally, IM or IV and co, give Epinephrine 0.3mg/ IV at a rate of 100-150ml/ 66 pounds: Diphenhydra	nfusion. ontact physician 0.3ml subcutaneously, hr and contact physician. mine 1.25mg/kg orally, IM
Is this the first dose? ☐ Yes ☐ No If no, list product ADMINISTER IVIG: ☐ 2 grams/kg over days, as a loading dose, then gm/kg or grams ☐ Rate per manufacturer recommendation ☐ Other Other ☐ Other PRE-MEDICATIONS ☐ Diphenhydramine (Benadryl) 25-50 mg orally before infusion ☐ Acetaminophen (Tylenol) 325-650 mg orally before infusion ☐ Other ☐ Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS (Adults or Children greater than 66 pounds or 30 kg: • For mild reaction: give Diphenhydramine 50 mg orally, IM or IV • For moderate reaction: stop infusion, give Diphenhydramine 50m • For Severe reaction w/breathing problem: stop infusion, call 911 Diphenhydramine 50 mg IV or IM. Begin NS 0.9% NaCI 500ml Note: Dosage adjustment necessary for children less than 30kg or or IV with a maximum of 50mg. If Epinephrine is needed 0.15mg/0	Date of last infusion	of for cycle(s) cycle(s) on fusion. on tact physician 0.3ml subcutaneously, thr and contact physician. mine 1.25mg/kg orally, IM usly
Is this the first dose? Yes No If no, list product ADMINISTER IVIG: 2 grams/kg over days, as a loading dose, then gm/kg or grams Rate per manufacturer recommendation Other Other PRE-MEDICATIONS Diphenhydramine (Benadryl) 25-50 mg orally before infusion Acetaminophen (Tylenol) 325-650 mg orally before infusion Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS of Adults or Children greater than 66 pounds or 30 kg: For mild reaction: give Diphenhydramine 50 mg orally, IM or IV For moderate reaction: stop infusion, give Diphenhydramine 50m For Severe reaction w/breathing problem: stop infusion, call 911 Diphenhydramine 50 mg IV or IM. Begin NS 0.9% NaCI 500ml Note: Dosage adjustment necessary for children less than 30kg or or IV with a maximum of 50mg. If Epinephrine is needed 0.15mg/0 Nursing: Start PIV as required for administration and nurse to administration and nurse to administration.	Date of last infusion	nfusion. ontact physician 0.3ml subcutaneously, hr and contact physician. mine 1.25mg/kg orally, IM asly Access: Peripheral PICC
Is this the first dose? Yes No If no, list product ADMINISTER IVIG: 2 grams/kg over days, as a loading dose, then gm/kg or grams Rate per manufacturer recommendation Other Other PRE-MEDICATIONS Diphenhydramine (Benadryl) 25-50 mg orally before infusion Acetaminophen (Tylenol) 325-650 mg orally before infusion Other ADVERSE/ANAPHYLACTIC REACTIONS: PER INS STANDARDS Of Adults or Children greater than 66 pounds or 30 kg: For mild reaction: give Diphenhydramine 50 mg orally, IM or IV For moderate reaction: stop infusion, give Diphenhydramine 50m orally Diphenhydramine 50mg IV or IM. Begin NS 0.9% NaCI 500ml Note: Dosage adjustment necessary for children less than 30kg or or IV with a maximum of 50mg. If Epinephrine is needed 0.15mg/0	Date of last infusion	of for cycle(s) cycle(s) on fusion. on tact physician 0.3ml subcutaneously, thr and contact physician. mine 1.25mg/kg orally, IM usly

By signing this form and utilizing our services, you are authorizing the pharmacy and it's employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.